

Observational constraints on spatial anisotropy of G from orbital motions

Lorenzo Iorio¹

Ministero dell'Istruzione, dell'Università e della Ricerca (M.I.U.R.)-Istruzione

Fellow of the Royal Astronomical Society (F.R.A.S.)

International Institute for Theoretical Physics and High Mathematics Einstein-Galilei

Permanent address for correspondence: Viale Unità di Italia 68, 70125, Bari (BA), Italy

`lorenzo.iorio@libero.it`

Received _____; accepted _____

ABSTRACT

A phenomenological anisotropic variation $\Delta G/G$ of the Newtonian gravitational coupling parameter G , if real, would affect the orbital dynamics of a two-body gravitationally bound system in a specific way. We analytically work out the long-term effects that such a putative modification of the usual Newtonian inverse-square law would induce on the trajectory of a test particle orbiting a central mass. Without making any a-priori simplifying assumptions concerning the orbital configuration of the test particle, it turns out that its osculating semi-major axis a , eccentricity e , pericenter ϖ and mean anomaly $\mathcal{M}$ undergo long-term temporal variations, while the inclination I and the node Ω are left unaffected. Moreover, the radial and the transverse components of the position and the velocity vectors $\boldsymbol{r}$ and $\boldsymbol{v}$ of the test particle experience non-vanishing changes per orbit, contrary to the out-of-plane ones. Then, we compute our theoretical predictions for some of the major bodies of the solar system by orienting the gradient of $G(\boldsymbol{r})$ towards the Galactic Center and keeping it fixed over the characteristic timescales involved. By comparing our calculation to the latest observational determinations for the same bodies, we infer $\Delta G/G \leq 10^{-17}$ over about 1 au. Finally, we consider also the Supermassive Black Hole hosted by the Galactic Center in Sgr A* and the main sequence star S2 orbiting it in about 16 yr, obtaining just $\Delta G/G \leq 10^{-2}$ over 1 kau.

Subject headings: Experimental studies of gravity; Experimental tests of gravitational theories; Modified theories of gravity; Orbital and rotational dynamics

PACS: 04.80.-y; 04.80.Cc; 04.50.Kd; 96.25.De

1. Introduction

The possibility that the Newtonian coupling parameter G may experience macroscopic spacetime variations ranging from laboratory to cosmological scales has been investigated both theoretically (Sciama 1953; Brans & Dicke 1961; Dicke 1962; Long 1974; Gershteyn & Gershteyn 1988; Linde 1990; De Sabbata et al. 1992; Melnikov 1994a,b; Capozziello et al. 1996; Melnikov 1996; Capozziello & de Ritis 1997; Capozziello et al. 1998; Drinkwater et al. 1998; Fischbach & Talmadge 1998; Barrow & O’Toole 2001; Danielsson 2001; Krause & Fischbach 2001; Murphy et al. 2001; Uzan 2003; de Sabbata et al. 2004; Capozziello 2005; Clifton et al. 2005; Hamber & Williams 2005; Bailey & Kostelecký 2006; Brownstein & Moffat 2006; García-Berro et al. 2007; Bertolami et al. 2008; Capozziello & Francaviglia 2008; Adelberger et al. 2009; Uzan 2009; Rami 2010; Kostelecký & Tasson 2011; Uzan 2011) and experimentally/observationally (Wagoner 1970; Vinti 1972; Ulrich 1974; Long 1976; Warburton & Goodkind 1976; Mikkelsen & Newman 1977; Anderson et al. 1978; Blinnikov 1978; Hut 1981; Chan et al. 1982; Kislik 1983; Chan & Paik 1984; Gillies 1987; Burgess & Cloutier 1988; Talmadge et al. 1988; Krauss & White 1992; Izmailov et al. 1993; Paik et al. 1994; Bertolami & Garcia-Bellido 1996; Gillies 1997; Gaztañaga et al. 2002; Gershteyn et al. 2002, 2004; Unnikrishnan & Gillies 2002a,b; Adelberger et al. 2003; Long 2003; Abramyan 2004; Barrow 2005; Boucher 2005; Kononogov & Mel’nikov 2005; García-Berro et al. 2007; Iorio 2007; Bertolami & Santos 2009; Newman et al. 2009; Li 2009; Lämmerzahl 2011; Piedipalumbo et al. 2011; Uzan 2011) since the early insights by Milne (1935, 1937), Dirac (1937) and Jordan (1937, 1939).

In this paper we deal with possible smooth anisotropic spatial variations of G , i.e. we consider the case $G = G(\mathbf{r})$ from a purely phenomenological point of view. We stress that in our analysis we do not rely upon any specific theoretical scheme encompassing such a spatial variability of G : the interested reader may want to consult the previously cited specialized literature. Quite generally, we express a putative anisotropic dependence of G on the spatial coordinates by parameterizing it with a gradient ∇G along a fixed direction in space $\hat{\xi}$ as

$$G(\mathbf{r}) \simeq G_0 + \nabla G_0 \cdot \mathbf{r}, \quad (1)$$

with

$$\nabla G_0 = |\nabla G_0| \hat{\xi}. \quad (2)$$

The subscript “0” in eq. (1) and eq. (2) refers to quantities evaluated at the origin of the spatial coordinates which, in our case, coincides with a generic body of mass M acting as source of the gravitational field. By means of eq. (1) we are assuming that the putative variations of G are rather smooth over the spatial extensions considered. A change of G

like that of eq. (1) is usually absent in the standard alternative metric theories of gravity treated within the Parameterized Post-Newtonian (PPN) framework (Will 1993) in which G may depend on the velocity $\mathbf{V}$ of the frame in which the experiments are performed with respect to a preferred frame (Will 1971); see, e.g., Vinti (1972), Damour & Esposito-Farèse (1994) and Will (1993) for analyses of the orbital consequences of such kind of anisotropies. A spatial anisotropy of G depending on the angle between the line of interaction of two gravitating bodies and a reference direction with respect to distant stars was experimentally investigated by¹ Gershteyn et al. (2002, 2004) in a series of Earth-based laboratory investigations which were subsequently critically analyzed by Unnikrishnan & Gillies (2002a). Unnikrishnan & Gillies (2002a) remarked that, in general, spatial anisotropies of G may occur depending on how nearby masses and their distribution can affect the gravitational interaction between two bodies, and also because of preferred frame effects. If the modification of the gravitational interaction depends on the gravitational potential generated by other masses (Brans & Dicke 1961; Sciama 1953) in a somewhat Machian fashion, then the most distant ones dominate and the spatial anisotropy is expected to be small (Unnikrishnan & Gillies 2002a). It is, then, reasonable to expect that the Galaxy yields the most important contribution to the anisotropy by assuming $\hat{\boldsymbol{\xi}}$ directed towards the Galactic Center (GC) (Unnikrishnan & Gillies 2002a).

In the framework of our parameterisation of eq. (1) and eq. (2), such a scenario may offer, in principle, interesting observational perspectives if suitable astronomical bodies are chosen. Indeed, if we insert eq. (1) in the usual expression of the Newtonian inverse-square law, a small modification of it occurs

$$\mathbf{A} = -\frac{M(\nabla G_0 \cdot \mathbf{r})\mathbf{r}}{r^3} = -\frac{M|\nabla G_0|(\hat{\boldsymbol{\xi}} \cdot \mathbf{r})\mathbf{r}}{r^3}. \quad (3)$$

Now, the ecliptic coordinates of the GC are (Reid & Brunthaler 2004)

$$\begin{aligned} \lambda_{\text{GC}} &= 183.15 \text{ deg}, \\ \beta_{\text{GC}} &= -5.61 \text{ deg}, \end{aligned} \quad (4)$$

so that the angle between $\hat{\boldsymbol{\xi}}$ and $\mathbf{r}$ for, say, a typical planet of the solar system is rather small. Thus,

$$\frac{\Delta G}{G} \sim \frac{|\nabla G_0| r}{G_0}. \quad (5)$$

Unnikrishnan & Gillies (2002a) argued that the order of magnitude of the Galactic-induced anisotropy is

$$\frac{\Delta G}{G} \sim \delta = \frac{GM_{\text{Gal}}}{c^2 d} \sim 10^{-6}, \quad (6)$$

¹They claimed to have measured an anisotropy as large as 5.4×10^{-4} with a torsion balance.

where c is the speed of light in vacuum, $M_{\text{Gal}} \sim 10^{12} M_{\odot}$ (Battaglia et al. 2005) is the mass of the Galaxy, and $d = 8.28$ kpc (Gillessen et al. 2009) is the distance from the GC. In our picture, it would naively be equivalent to perturbing accelerations

$$A \lesssim 10^{-6} A_{\text{N}}, \quad (7)$$

where the standard inverse-square Newtonian accelerations A_{N} are

$$4 \times 10^{-2} \text{ m s}^{-2} \leq A_{\text{N}} \leq 4 \times 10^{-6} \text{ m s}^{-2} \quad (8)$$

for the major bodies of the solar system. In principle, perturbing accelerations as large as

$$A \sim 10^{-8} - 10^{-12} \text{ m s}^{-2} \quad (9)$$

may have interesting observational consequences. We will analytically work out them in detail. Indeed, relying upon simple order-of-magnitude evaluations may be misleading since important factors of the order of $\mathcal{O}(e^j)$, $j = 1, 2, \dots$ or $\mathcal{O}(e^{-j})$, $j = 1, 2, \dots$ in the usually small eccentricities e of the bodies adopted as probes may be neglected. Unnikrishnan & Gillies (2002a) performed a preliminary calculation concerning the Earth-Moon system. They started from a certain value $|\dot{G}/G| \leq 4 \times 10^{-12} \text{ yr}^{-1}$ (Dickey et al. 1994) of the upper bound in the fractional time change of G obtained with the Lunar Laser ranging (LLR) technique. Then, Unnikrishnan & Gillies (2002a) stated that the same analysis can be useful as far as the spatial anisotropy of G is concerned as well. They noticed that an anisotropic spatial variation of G like that of eq. (1) should exhibit a harmonic signal with the same approximate monthly periodicity of the orbital lunar motion as the line joining the Earth and the Moon sweeps out different directions with respect to the GC direction. Finally, since the accuracy with which it is possible to measure a periodic signal may be of the same order of, or better than that for a secular trend, Unnikrishnan & Gillies (2002a) concluded by inferring²

$$\frac{\Delta G}{G} \leq 4 \times 10^{-12}. \quad (10)$$

Repeating the same reasonings with the latest results from LLR (Williams et al. 2004; Müller & Biskupek 2007; Williams et al. 2009; Hofmann et al. 2010) would yield

$$\frac{\Delta G}{G} \leq (1 - 0.4) \times 10^{-12}, \quad (11)$$

in neat disagreement the results by Gershteyn et al. (2002). Gershteyn et al. (2002) pointed out that such a disagreement exists if it is assumed that the G anisotropy depends neither on the magnitude of the interacting masses nor on the distance between them;

²It is so also because 1 year contains almost 12 lunar cycles.

Gershteyn et al. (2002) remarked that the masses and the distances involved in the analysis by Unnikrishnan & Gillies (2002a) drastically differ from those used by Gershteyn et al. (2002) in their experiment.

We propose to obtain much more accurate bounds than that in eq. (10) by calculating in detail all the orbital effects of eq. (3) which, actually, depends on the mutual distance between M and the test body: we will assume that it is independent of their masses. As far as other performed and/or proposed astronomical tests of $G(\mathbf{r})$ are concerned (Wagoner 1970; Vinti 1972; Ulrich 1974; Warburton & Goodkind 1976; Mikkelsen & Newman 1977; Anderson et al. 1978; Blinnikov 1978; Hut 1981; Kislik 1983; Burgess & Cloutier 1988; Talmadge et al. 1988; Abramyan 2004; Iorio 2007; Bertolami & Santos 2009; Li 2009), quite different explicit theoretical isotropic models and empirical approaches have been followed so far; in particular, an exponential Yukawa-type model and the third Kepler law have often been adopted.

The paper is organized as follows. In Section 2 we analytically work out various orbital effects caused by a phenomenological dipole-type spatial variation of G like that of eq. (1) averaged over one orbital revolution of a test particle. In Section 3 a comparison with the latest solar system planetary observations is made; we also consider the stellar system around the Galactic black hole. Section 4 summarizes our findings and contain the conclusions.

2. Calculation of the orbital effects

The orbital motions of test particles are, in principle, affected by eq. (3) whose effects can be worked out with standard perturbative techniques (Brouwer & Clemence 1961; Bertotti et al. 2003). By defining³

$$\psi_0 \doteq \frac{|\nabla G_0|}{G_0}, \quad \mu_0 \doteq G_0 M, \quad (12)$$

the radial component A_R of eq. (3), evaluated onto the unperturbed Keplerian ellipse, is

$$A_R = -\frac{\psi_0 \mu_0 (1 + e \cos f) \left\{ \cos u \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \sin u \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}}{a (1 - e^2)}, \quad (13)$$

the transverse and out-of-plane components A_T, A_N vanish. In eq. (13) $a, I, \Omega, u \doteq \omega + f, \omega, f$ are the osculating semi-major axis, the inclination of the orbital plane to the reference

³Notice that $[\psi_0] = \text{L}^{-1}$, while $[G_0 M] = \text{L}^3 \text{T}^{-2}$, as usual.

$\{x, y\}$ plane, the longitude of the ascending node⁴, the argument of latitude, the argument of pericentre⁵ and the true anomaly⁶, respectively, of the test particle. Notice that eq. (13) clearly shows that the putative anisotropic G effect depends, among other things, on the distance between the two bodies as well. By assuming $\hat{\xi}$ constant over one orbital revolution of the test particle, a straightforward first-order application of the Gauss perturbative equations (Brouwer & Clemence 1961; Bertotti et al. 2003) yields for a , e , the longitude⁷ of the pericenter $\varpi \doteq \Omega + \omega$, and $\mathcal{M}$ the following non-vanishing long-term temporal rates of change

$$\begin{aligned}\frac{da}{dt} &= -\frac{2\psi_0\mu_0}{aen} \left\{ -\sin \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \cos \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \frac{de}{dt} &= -\frac{\psi_0\mu_0(1-e^2)}{a^2e^2n} \left\{ -\sin \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \cos \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \frac{d\varpi}{dt} &= -\frac{\psi_0\mu_0(1-e^2)}{a^2e^3n} \left\{ \cos \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \sin \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \frac{d\mathcal{M}}{dt} &= \frac{\psi_0\mu_0(1-e^2)^{3/2}}{a^2e^3n} \left\{ \cos \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \sin \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}.\end{aligned}\tag{14}$$

The inclination I and the node Ω are left unaffected. Notice that the quantity n entering eq. (14) is the unperturbed Keplerian mean motion, i.e. $n \doteq \sqrt{\mu_0/a^3}$. The long-term rates of change of eq. (14) are exact in the sense that no a-priori assumptions concerning $\hat{\xi}$ and the orbital configuration of the test particle were made. All the formulas of eq. (14) are singular for $e \rightarrow 0$; however, it is just an unphysical artifact which can be cured by adopting the well-known non-singular elements (Brouwer & Clemence 1961; Broucke & Cefola 1972)

$$\begin{aligned}h &\doteq e \sin \varpi, \\ k &\doteq e \cos \varpi, \\ l &\doteq \varpi + \mathcal{M}.\end{aligned}\tag{15}$$

⁴It is an angle in the reference $\{xy\}$ plane counted from the reference x direction to the line of the nodes, which is the intersection of the orbital plane with the reference $\{xy\}$ plane.

⁵It is an angle in the orbital plane reckoning the point of closest approach with respect to the line of the nodes.

⁶It is a time-dependent angle in the orbital plane determining the instantaneous position of the test particle with respect to the pericentre.

⁷It is a “dogleg” angle.

It is also important to remark that the validity of eq. (14) is not restricted to any specific reference frame, being, instead, quite general.

It is useful to work out the radial, transverse, and out-of-plane shifts over one orbital revolution of the position and velocity vectors $\mathbf{r}$ and $\mathbf{v}$ as well. They can be analytically worked out according to Casotto (1993). For the shifts $\Delta R, \Delta T, \Delta N$ of $\mathbf{r}$ we have

$$\begin{aligned}\Delta R &= \frac{2\pi\psi_0 a^2 (1-e^2)^2}{e^2} \left\{ -\sin \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \cos \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \Delta T &= \frac{4\pi\psi_0 a^2 (1-e^2)}{e^2} \left\{ \cos \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \sin \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \Delta N &= 0.\end{aligned}\tag{16}$$

The shifts $\Delta v_R, \Delta v_T, \Delta v_N$ of $\mathbf{v}$ are

$$\begin{aligned}\Delta v_R &= -\frac{2\pi\psi_0 a^2 n (1+e)^{3/2}}{e^2 \sqrt{1-e}} \left\{ \cos \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \sin \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \Delta v_T &= -\frac{2\pi\psi_0 a^2 n \sqrt{1-e^2}}{e^2} \left\{ -\sin \omega \left(\hat{\xi}_x \cos \Omega + \hat{\xi}_y \sin \Omega \right) + \cos \omega \left[\hat{\xi}_z \sin I + \cos I \left(\hat{\xi}_y \cos \Omega - \hat{\xi}_x \sin \Omega \right) \right] \right\}, \\ \Delta v_N &= 0.\end{aligned}\tag{17}$$

Concerning the singularities occurring for $e \rightarrow 0$, also for eq. (16) and eq. (17) the same considerations as for eq. (14) hold.

3. Confrontation with the observations

Here we compare our theoretical results of Section 2 with the latest observationally determined quantities pertaining the orbital motions of some of the major bodies of the solar system obtained by processing long data records of various types with different ephemerides (Pitjeva 2007; Fienga et al. 2011, 2010; Pitjeva 2010) in order to preliminarily infer upper bounds on ψ_0 . In principle, one should explicitly include eq. (1) in the dynamical force models fit to the observations and re-process the entire data set with such an *ad-hoc* modified theory by varying the parameters to be estimated, the data, etc. Otherwise, the putative signal may be partly or totally absorbed in the estimation of, say, the initial state vectors. However, this is beyond the scopes of our work.

In computing the anomalous effects for different bodies, we refer their orbital configurations to a heliocentric frame with mean ecliptic and equinox at the epoch J2000.

In it the unit vector pointing to the GC is

$$\begin{aligned}\hat{\xi}_x &= -0.993, \\ \hat{\xi}_y &= -0.054, \\ \hat{\xi}_z &= -0.097.\end{aligned}\tag{18}$$

In Table 1 we quote reasonable evaluations for the secular variations of the semi-major axes a and the eccentricities e of the inner planets of the solar system for which the most accurate data are currently available. They were computed by dividing the formal, statistical $1 - \sigma$ errors in a, e of the EPM2006 ephemerides Pitjeva (2007) by the time interval $\Delta t = 93$ yr covered by the observations used for constructing them. The realistic uncertainties may

Table 1: Formal uncertainties in the secular variations of the semi-major axes a and the eccentricities e of the inner planets of the solar system. They were obtained by dividing the formal errors in a and e in Table 3 of Pitjeva (2007) by the time interval $\Delta t = 93$ yr (1913-2006) of the data records used for the EPM2006 ephemerides (Pitjeva 2007). The errors in e were computed as $\sigma_e = \sqrt{\left(\frac{\partial e}{\partial h}\right)^2 \sigma_h^2 + \left(\frac{\partial e}{\partial k}\right)^2 \sigma_k^2}$. The realistic uncertainties may be up to one order of magnitude larger.

Planet	σ_a (m yr ⁻¹)	σ_e (yr ⁻¹)
Mercury	3×10^{-3}	4.43×10^{-12}
Venus	2×10^{-3}	1.8×10^{-13}
Earth	1×10^{-3}	5×10^{-14}
Mars	3×10^{-3}	5×10^{-14}

be up to one order of magnitude larger. Table 2 displays the latest determinations of the corrections $\Delta\dot{\varpi}$ to the standard Newtonian/Einsteinian secular precessions of the perihelia of the planets of the solar system recently obtained with different ephemerides (Fienga et al. 2011, 2010; Pitjeva 2010). Since ψ_0 enters eq. (14) as a free, adjustable parameter, we select the values for it which make the putative anomalous effects of eq. (14) not larger than the empirically obtained bounds in Table 1 and Table 2. As a result, we are able to obtain reasonable guesses concerning $\Delta G/G$ at different heliocentric distances by posing $\Delta G \sim \sigma_{\psi_0} \langle r \rangle$. The results are shown in Table 3. Even by re-scaling the bounds obtained from Table 1 by one order of magnitude, the anisotropy of G in the planetary regions of the solar system is very tightly constrained, being of the order of $10^{-15} - 10^{-17}$.

It may be interesting to consider a completely different astronomical scenario, both from the point of view of its components and of the distance scales involved. In Table 4 we quote the relevant physical and orbital parameters of the system constituted by the

Table 2: Estimated corrections $\Delta\dot{\varpi}$, in milliarcseconds per century (mas cty^{-1}), to the standard Newtonian-Einsteinian secular precessions of the longitudes of the perihelia ϖ of the eight planets of the solar system plus Pluto determined with the EPM2008 (Pitjeva 2010), the INPOP08 (Fienga et al. 2010), and the INPOP10a (Fienga et al. 2011) ephemerides. Only the usual Newtonian-Einsteinian dynamics was modelled, so that, in principle, the corrections $\Delta\dot{\varpi}$ may account for any other unmodelled/mismodelled dynamical effect. Concerning the values quoted in the third column from the left, they correspond to the smallest uncertainties reported by Fienga et al. (2010). Note the small uncertainty in the correction to the precession of the terrestrial perihelion, obtained by processing Jupiter VLBI data (Fienga et al. 2010).

Planet	$\Delta\dot{\varpi}$ (Pitjeva 2010)	$\Delta\dot{\varpi}$ (Fienga et al. 2010)	$\Delta\dot{\varpi}$ (Fienga et al. 2011)
Mercury	-4 ± 5	-10 ± 30	0.2 ± 3
Venus	24 ± 33	-4 ± 6	–
Earth	6 ± 7	0 ± 0.016	–
Mars	-7 ± 7	0 ± 0.2	–
Jupiter	67 ± 93	142 ± 156	–
Saturn	-10 ± 15	-10 ± 8	0 ± 2
Uranus	-3890 ± 3900	0 ± 20000	–
Neptune	-4440 ± 5400	0 ± 20000	–
Pluto	2840 ± 4510	–	–

Table 3: Upper bounds on the anisotropic percent variation $\Delta G/G$ inferred from $\dot{a}, \dot{e}, \dot{\varpi}$ for the inner planets of the solar system. We posed $\Delta G \lesssim \sigma_{\psi_0} \langle r \rangle = \sigma_{\psi_0} a (1 + e^2/2)$ for each planet, where σ_{ψ_0} was obtained by comparing the theoretical predictions of eq. (14) to the uncertainties listed in Table 1 and Table 2.

Planet	$\frac{\Delta G}{G} \big _{\dot{a}}$	$\frac{\Delta G}{G} \big _{\dot{e}}$	$\frac{\Delta G}{G} \big _{\dot{\varpi}}$
Mercury	2×10^{-16}	8×10^{-15}	3×10^{-13}
Venus	9×10^{-18}	1×10^{-18}	1×10^{-17}
Earth	1×10^{-17}	2×10^{-18}	3×10^{-18}
Mars	4×10^{-16}	3×10^{-16}	3×10^{-15}

Supermassive Black Hole (SBH) hosted by the GC in Sgr A* and the main sequence star S2 orbiting it in about 16 yr at a distance of approximately 1 kau (Gillessen et al. 2009). In this case, the angular elements refer to a coordinate system whose reference z axis is directed along the line of sight: the reference $\{x, y\}$ plane coincides with the plane of the

sky⁸, with the x axis pointing towards the Celestial North Pole. The results of eq. (14)

Table 4: Relevant physical and orbital parameters of the SBH-S2 system in Sgr A*. The Keplerian orbital elements of S2 were retrieved from Table 1 of (Gillessen et al. 2009). The figure for the gravitational parameter μ_0 comes from a multi-star fit yielding $\mu_0 = 4.30 \times 10^6 \mu_\odot$ (Gillessen et al. 2009): it yields a Schwarzschild radius as large as $r_g = 0.084$ au. The quoted value in m for the semi-major axis of S2 was obtained by multiplying its angular value $a = 0.1246$ arcsec (Gillessen et al. 2009) by the distance of the SBH $d = 8.28$ kpc (Gillessen et al. 2009): it corresponds to 1031.69 au, so that the orbital period of S2 is $P_b = 15.98$ yr. The uncertainty σ_ω in the secular precession of S2 can naively be obtained by dividing the error in ω , quoted in Table 1 of (Gillessen et al. 2009), by the time interval ΔT covered by the observations used which is almost equal to P_b .

μ_0 ($\text{m}^3 \text{s}^{-2}$)	a (m)	e	I (deg)	Ω (deg)	ω (deg)	σ_ω (arcsec yr ⁻¹)
5.70×10^{26}	1.54×10^{14}	0.8831	134.87	226.53	64.98	182

are applicable to S2 as well since its mass is about five orders of magnitude smaller than that of the SBH: clearly, in this case it is $\hat{\xi} = -\hat{z}$. Concerning the use of eq. (14) in the Sgr A* scenario, one may wonder why an essentially Newtonian scheme is adopted instead of a general relativistic one. In principle, one could assume as reference path a fully post-Newtonian one⁹ (Calura et al. 1997, 1998), and work out the effects of a given small extra-acceleration like eq. (3) with respect to it according to the perturbative scheme set up by Calura et al. (1997, 1998), which is a general relativistic generalization of another standard perturbative approach based on the planetary Lagrange equations (Bertotti et al. 2003). Actually, it is, in practice, useless since the only addition with respect to the orbital effects like, e.g., the precession of the pericenter, resulting from the standard scenario would consist of further, small mixed GTR-perturbation orbital effects, completely irrelevant in strengthening the bounds inferred. Viewed from a different point of view, the ratio of the average distance $\bar{r}_{\text{S2}}$ of S2 from the SBH to its Schwarzschild radius r_g is, after all, as large as 1.7×10^4 .

By using the expression for the putative precession of the stellar pericenter in eq. (14), it can be compared to the present-day uncertainty in observationally determining its secular rate in Table 4. In this case, the constraints on the G anisotropy are very weak. Indeed, we

⁸It is a plane tangent to the Celestial Sphere at the point where the object of interest is located.

⁹In doing so, it would be implicitly assumed that the effects due to a putative G anisotropy are smaller than the post-Newtonian ones as well.

have just

$$\psi_0^{-1} \geq 0.45 \text{ pc}, \quad (19)$$

corresponding to

$$\frac{\Delta G}{G} \leq 1.5 \times 10^{-2} \quad (20)$$

over about 1 kau. Note that the bound of eq. (20) is tighter by one order of magnitude than that could be inferred by simply posing

$$\frac{\Delta G}{G} \lesssim \frac{\sigma_\mu}{\mu_0} = 1.2 \times 10^{-1}, \quad (21)$$

from $\sigma_\mu = 0.50 \times 10^6 \mu_\odot$ (Gillessen et al. 2009).

4. Summary and conclusions

We looked at phenomenological anisotropic spatial variations $\Delta G/G$ of the Newtonian gravitational coupling parameter G , in the form $\Delta G = \nabla G \cdot \mathbf{r}$, and analytically worked out the impact that they may have on the trajectory of a test particle orbiting a central body of mass M . More specifically, we focussed on the cumulative orbital changes obtained perturbatively by averaging over one period of revolution of the test particle the effects due to $\Delta G/G$ on its path. As a result, the osculating semi-major axis a , the eccentricity e , the pericenter ϖ and the mean anomaly $\mathcal{M}$ of the orbiter experience non-vanishing long-term changes which depend on the overall orbital geometry of the test particle and on the direction $\hat{\xi}$ of the putative gradient ∇G . We analytically worked out the long-term variations per orbit $\Delta R, \Delta T, \Delta N$ and $\Delta v_R, \Delta v_T, \Delta v_N$ of the position and velocity vectors $\mathbf{r}$ and $\mathbf{v}$ of the test particle as well. We found that both the radial and the transverse components of $\mathbf{r}$ and $\mathbf{v}$ are affected by long-term changes per orbit, while the out-of-plane ones are left unaffected. We kept $\hat{\xi}$ fixed during the integrations: moreover, no a-priori simplifying assumptions on e and I were assumed, so that our results are exact in this respect.

Then, we compared our theoretical predictions to the most recently determined observational quantities for some of the major bodies of the solar system. By assuming that the dominant contribution to the hypothetical anisotropy of G is due to the Galaxy, we took $\hat{\xi}$ directed towards the Galactic Center, which has a small inclination with respect to the ecliptic. By using the heliocentric orbits of the inner planets we were able to constrain $\Delta G/G$ to a level of about 10^{-17} over ~ 1 au, several orders of magnitude better than in previous analyses based on Lunar Laser Ranging only. We looked also at the star S2 orbiting the Galactic black hole at a distance of about 1 kau along a highly elliptical ellipse, but, in this case, we got just $\Delta G/G \lesssim 10^{-2}$.

REFERENCES

- Abramyan G.L., 2004, On The Spatial Variation of the Gravitational Constant,
<http://www.agl-planets.narod.ru/pdf1/engge.pdf>
- Adelberger E. G., Heckel B. R., Nelson A. E., 2003, Annual Review of Nuclear and Particle Science, 53, 77
- Adelberger E. G., Gundlach J. H., Heckel B. R., Hoedl S., Schlamminger S., 2009, Progress in Particle and Nuclear Physics, 62, 102
- Anderson J. D., Keesey M. S. W., Lau E. L., Standish E. M., Newhall X. X., 1978, Acta Astronautica, 5, 43
- Bailey Q. G., Kostelecký V. A., 2006, Phys. Rev. D, 74, 045001
- Barrow J. D., O’Toole C., 2001, Mon. Not. Roy. Astron. Soc., 322, 585
- Barrow J. D., 2005, Phys. Rev. D, 71, 083520
- Battaglia G., Helmi A., Morrison H., Harding P., Olszewski E. W., Mateo M., Freeman K. C., Norris J., Shectman S. A., 2005, Mon. Not. Roy. Astron. Soc., 364, 433
- Bertolami O., Garcia-Bellido J., 1996, Int. J. Mod. Phys. D, 5, 363
- Bertolami O., Páramos J., Turyshev S.G., 2008, General Theory of Relativity: Will it survive the Next Decade?. In: Dittus H., Lämmerzahl C., Turyshev S. G. (eds.) Lasers, Clocks and Drag-Free Control. Exploration of Relativistic Gravity in Space. Astrophysics and Space Science Library Volume 349, Springer, Berlin, pp. 27-74
- Bertolami O., Santos N. M. C., 2009, Phys. Rev. D, 79, 127702
- Bertotti B., Farinella P., Vokrouhlický D., 2003, Physics of the Solar System, Kluwer Academic Press, Dordrecht
- Blinnikov S. I., 1978, Astrophys. Space Sci., 59, 13
- Boucher V., 2005, Gravitation and Cosmology, 11, 71
- Brans C., Dicke R. H., 1961, Phys. Rev., 124, 925
- Broucke R. A., Cefola P. J., 1972, Celest. Mech. Dyn. Astron., 5, 303
- Brouwer D., Clemence G., 1961, Methods of Celestial Mechanics, New York, Academic Press.
- Brownstein J. R., Moffat J. W., 2006, Class. Quantum Grav., 23, 3427

- Burgess C. P., Cloutier J., 1988, *Phys. Rev. D*, 38, 2944
- Calura M., Fortini P., Montanari E., 1997, *Phys. Rev. D*, 56, 4782
- Calura M., Fortini P., Montanari E., 1998, *Class. Quantum Grav.*, 15, 3121
- Capozziello S., de Ritis R., Rubano C., Scudellaro P., 1996, *Riv. N. Cimento*, 19, 2
- Capozziello S., de Ritis R., 1997, *Gen. Rel. Grav.*, 29, 1425
- Capozziello S., de Ritis R., Marino A. A., 1998, *Gen. Rel. Grav.*, 30, 1247
- Capozziello S., 2005, Newtonian Limit of Extended Theories of Gravity. In: Reimer A. (ed.) *Quantum Gravity Research Trends. Horizons in World Physics*, Volume 250, NOVA, Hauppauge, pp. 223-272
- Capozziello S., Francaviglia M., 2008, *Gen. Relativ. Gravit.*, 40, 357
- Casotto S., 1993, *Celest. Mech. Dyn. Astron.*, 55, 209
- Chan H. A., Moody M. V., Paik H. J., 1982, *Phys. Rev. Lett.*, 49, 1745
- Chan H. A., Paik H. J., 1984, in Taylor B. N., Phillips W. D. (eds.) *Precision Measurement and Fundamental Constants II*, National Bureau of Standards Special Publication 617, Gaithersburg, Maryland, p. 601
- Clifton T., Mota D. F., Barrow J. D., 2005, *Mon. Not. Roy. Astron. Soc.*, 358, 601
- Damour T., Esposito-Farèse G., 1994, *Phys. Rev. D*, 49, 1693
- Danielsson U., 2001, *Rep. on Progr. in Phys.*, 64, 51
- de Sabbata V., Melnikov V. N., Pronin P. I., 1992, *Prog. Theor. Phys.*, 88, 623
- de Sabbata V., Gillies G. T., Melnikov V. N. (eds.), 2004, *The Gravitational Constant: Generalized Gravitational Theories and Experiments* *The Gravitational Constant: Generalized Gravitational Theories and Experiments*, Springer, Berlin
- Dicke R., 1962, *Phys. Rev.*, 126, 1875
- Dickey J. O., Bender P. L., Faller J. E., Newhall X. X., Ricklefs R. L., Ries J. G., Shelus P. J., Veillet C., Whipple A. L., Wiant J. R., Williams J. G., Yoder C. F., 1994, *Science*, 265, 482
- Dirac P. A. M., 1937, *Nature*, 139, 323
- Drinkwater M. J., Webb J. K., Barrow J. D., Flambaum V. V., 1998, *Mon. Not. Roy. Astron. Soc.*, 295, 457

- Fienga A., Laskar J., Kuchynka P., Leponcin-Lafitte C., Manche H., Gastineau M., 2010, Gravity tests with INPOP planetary ephemerides. In: Klioner S. A., Seidelmann P. K., Soffel M. H. (eds.) *Relativity in Fundamental Astronomy: Dynamics, Reference Frames, and Data Analysis*, Proceedings IAU Symposium No. 261, Cambridge University Press, Cambridge, pp. 159-169.
- Fienga A., Manche H., Kuchynka P., Laskar J., Gastineau M., 2011, Planetary and lunar ephemerides INPOP10A. In: Capitaine N. (ed.) *Proceedings of the Journées 2010 “Systèmes de Référence Spatio-Temporels”*, N. Capitaine (ed.), Observatoire de Paris, Paris, pp. 37-42.
- Fischbach E., Talmadge C., 1998, *The Search for Non Newtonian Gravity*. Springer, Berlin
- García-Berro E., Isern J., Kubyshev Y. A., 2007, *Astron. Astrophys. Rev.*, 14, 113
- Gaztañaga E., García-Berro E., Isern J., Bravo E., Domínguez I., 2002, *Phys. Rev. D*, 65, 023506
- Gershteyn M. L., Gershteyn L. I., 1988, A new approach to the electricity and gravity amalgamation, *Redaktsiya Izvestiya Vuzov Radiophysica*, Gorky, deposited in VINITI 28.09.88, No 7196-B88
- Gershteyn M. L., Gershteyn L. I., Gershteyn A. M., Karagioz O. V., 2002, *Gravitation and Cosmology*, 8, 243
- Gershteyn M. L., Gershteyn L. I., Gershteyn A., Karagioz O.V., 2004, *Infinite Energy*, 10, 26
- Gillessen S., Eisenhauer F., Fritz T.K., Bartko H., Dodds-Eden K., Pfuhl O., Ott T., Genzel R., 2009, *Astrophys. J.*, 707, L114
- Gillies G. T., 1987, *Metrologia*, 24, 44
- Gillies G. T., 1997, *Rep. Prog. Phys.*, 60, 151
- Hamber H. W., Williams R. M., 2005, *Phys. Rev. D*, 72, 044026
- Hofmann F., Müller J., Biskupek L., 2010, *Astron. Astrophys.*, 522, L5
- Hut P., 1981, *Phys. Lett. B*, 99, 174
- Iorio L., 2007, *Mon. Not. Roy. Astron. Soc.*, 376, 1727
- Izmailov V. P., Karagioz O.V., Kuznetsov V. A., Mel’nikov V. N., Roslyakov A. E., 1993, *Measurement Techniques*, 36, 1065
- Jordan P., 1937, *Naturwiss.*, 25, 513

- Jordan P., 1939, *Physik Z*, 113, 660
- Kislik M. D., 1983, *Sov. Astron. Lett.*, 9, 168
- Kononogov S. A., Mel’nikov V. N., 2005, *Measurement Techniques*, 48, 521
- Kostelecký V. A., Tasson J. D., 2011, *Phys. Rev. D*, 83, 016013
- Krause D. E., Fischbach E., 2001, in Lammerzahl C., Everitt C.W. F., Hehl F.W., eds, 2001, *Gyros, Clocks, Interferometers. . . : Testing Relativistic Gravity in Space*. Springer, Berlin, p. 292
- Krauss L. M., White M., 1992, *Astrophys. J.*, 397, 357
- Lämmerzahl C., 2011, Testing Basic Laws of Gravitation-Are Our Postulates on Dynamics and Gravitation Supported by Experimental Evidence? In: Blanchet L., Spallicci A., Whiting B. (eds.) *Mass and Motion in General Relativity*, Berlin, Springer, pp. 25-65
- Li L.-S., 2009, *Nuovo Cimento B*, 124, 849
- Linde A., 1990, *Phys. Lett. B*, 238, 160
- Long D. R., 1974, *Phys. Rev. D*, 9, 850
- Long D. R., 1976, *Nature*, 260, 417
- Long J. C., 2003, *Nature*, 421, 922
- Melnikov V. N., 1994a, *Int. J. Theor. Phys.*, 33, 1569
- Melnikov V. N., 1994b, Multidimensional classical and quantum cosmology and gravitation: exact solutions and variations of constants. In: Novello M. (ed.) *Cosmology and Gravitation*. Editions Frontières, Gif-sur-Yvette, pp. 147-244
- Melnikov V. N., 1996, Classical Solution in Multidimensional Cosmology. In: Novello M. (ed.) *Cosmology and Gravitation II*. Editions Frontières, Gif-sur-Yvette, p. 465
- Mikkelsen D. R., Newman M. J., 1977, *Phys. Rev. D*, 16, 919
- Milne E. A., 1935, *Relativity, gravitation and world structure*. Claredon Press, Oxford
- Milne E. A. , 1937, *Proc. Roy. Soc. A*, 158, 324
- Müller J., Biskupek L., 2007, *Class. Quantum Gravit.*, 24, 4533
- Murphy M. T., Webb J. K., Flambaum V. V., Dzuba V. A., Churchill C. W., Prochaska J. X., Barrow J. D., Wolfe A. M., 2001, *Mon. Not. Roy. Astron. Soc.*, 327, 1208

- Newman R. D., Berg E. C., Boynton P. E., 2009, *Space Sci. Rev.*, 148, 175
- Paik H. J., Moody M. V., 1994, *Class. Quantum Gravit.*, 11, A145
- Piedipalumbo E., Esposito G., Rubano C., Scudellaro P., 2011, arXiv:1104.3029
- Pitjeva E.V., 2007, Use of optical and radio astrometric observations of planets, satellites and spacecraft for ephemeris astronomy, in Jin W. J., Platais I., Perryman M. A. C. (eds.), *A Giant Step: from Milli- to Micro-arcsecond Astrometry Proceedings IAU Symposium No. 248*, Cambridge University Press, Cambridge, pp. 20-22
- Pitjeva E. V., 2010, EPM ephemerides and relativity. In: Klioner S. A., Seidelmann P. K., Soffel M. H., (eds.) *Relativity in Fundamental Astronomy: Dynamics, Reference Frames, and Data Analysis, Proceedings IAU Symposium No. 261*, Cambridge University Press, Cambridge, pp. 170-178.
- Rami El-N. A., 2010, *Brazilian J. Phys.*, 40, 92
- Reid M. J., Brunthaler A., 2004, *Astrophys. J.*, 616, 872
- Sciama D. W., 1953, *Mon. Not. Roy. Astron. Soc.*, 113, 34
- Talmadge C., Berthias J.-P., Hellings R. W., Standish E. M., 1988, *Phys. Rev. Lett.*, 61, 1159
- Ulrich R. K., 1974, *Astrophys. J.*, 188, 369
- Unnikrishnan C. S., Gillies G. T., 2002a, *Phys. Lett. A*, 305, 26
- Unnikrishnan C. S., Gillies G. T., 2002b, Constraints on gravitational screening. In: Edwards M. R. (ed.) *Pushing Gravity: New Perspectives on Le Sage's Theory of Gravitation*, Apeiron, Montreal, pp. 259-266
- Uzan J.-P., 2003, *Rev. Mod. Phys.*, 75, 403
- Uzan J.-P., 2009, *Space Sci. Rev.*, 148, 249
- Uzan J.-P., 2011, *Living Rev. Relativity*, 14, 2, URL (cited on 9 August 2011): <http://www.livingreviews.org/lrr-2011-2>
- Vinti J. P., 1972, *Celest. Mech. Dyn. Astron.*, 6, 198
- Wagoner R. V., 1970, *Phys. Rev. D*, 1, 3209
- Warburton R. J., Goodkind J. M., 1976, *Astrophys. J.*, 208, 881
- Will C.M., 1971, *Astrophys. J.*, 169, 141

Will C.M., 1993, Theory and experiment in gravitational physics. Revised edition, Cambridge University Press, Cambridge

Williams J. G., Turyshv S. G., Boggs D. H., 2004, Phys. Rev. Lett., 93, 261101

Williams J. G., Turyshv S. G., Boggs D. H., 2009, Int. J. Mod. Phys. D, 18, 1129